

# Widely separated binary systems of very low mass stars

N. PHAN-BAO<sup>1</sup>, E.L. MARTÍN<sup>2,3</sup>, C. REYLÉ<sup>4</sup>, T. FORVEILLE<sup>5,6</sup>, and J. LIM<sup>1</sup>

<sup>1</sup> Institute of Astronomy and Astrophysics, Academia Sinica, P.O. Box 23-141, Taipei 106, Taiwan, R.O.C.

<sup>2</sup> Instituto de Astrofísica de Canarias, C/ Vía Láctea s/n, E-38200 La Laguna (Tenerife), Spain.

<sup>3</sup> University of Central Florida, Dept. of Physics, PO Box 162385, Orlando, FL 32816-2385, USA.

<sup>4</sup> CNRS UMR6091, Observatoire de Besançon, BP1615, 25010 Besançon Cedex, France.

<sup>5</sup> Canada-France-Hawaii Telescope Corporation, 65-1238 Mamalahoa Highway, Kamuela, HI 96743 USA.

<sup>6</sup> Laboratoire d'Astrophysique de Grenoble, Université J. Fourier, B.P. 53, F-38041 Grenoble, France.

Received; accepted; published online

**Abstract.** In this paper we review some recent detections of wide binary brown dwarf systems and discuss them in the context of the multiplicity properties of very low-mass stars and brown dwarfs.

**Key words:** binary stars, very low mass stars, brown dwarfs, individual star: DENIS-P J0410-1251, LP 714-37

©0000 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

## 1. Introduction

Binary systems have been studied for decades to measure accurate stellar masses, and to test evolutionary models and star formation theories. Considerable attention has recently been paid to very low-mass (VLM) binaries in the solar neighborhood (Martín et al. 1999a; Close et al. 2002, 2003; Bouy et al. 2003; Burgasser et al. 2003; Forveille et al. 2005), as well as in nearby young open clusters and associations (Martín et al. 2003; Chauvin et al. 2004).


The properties of VLM binaries are an important constraint for models of star-formation and evolution. It has been debated in the literature whether the properties of VLM binaries and stellar binaries differ, implying different formation mechanisms (Kroupa et al. 2003), or whether the binary properties instead show continuous trends with decreasing primary mass, implying that VLM binaries form through the same processes as stellar binaries (Luhman 2004b). Clearly there is a need for a larger sample of observed VLM binaries, particularly at wide separations where few of them are known. One leading model of brown dwarf (BD) formation is that they form and are ejected in unstable multiple systems within small clusters. Since the ejection models (Reipurth & Clarke 2001; Bate et al. 2002) suggest that the binary BD systems that do exist must be close (separations  $\leq 10$  AU). The detection of wide VLM binary systems has thus become an important test of the ejection models. The first wide binary

BDs have been found in young ( $<10$  Myr) associations or clusters (Luhman 2004a; Chauvin et al. 2004). Very recently Phan-Bao et al. (2005) found a 33 AU ultracool binary system, Billères et al. (2005) have also reported a discovery of a 200 AU M8.0+L0 binary, both of them in the field and their detection based on the DENIS survey.

## 2. LP 714-37AB: the closest wide binary system of very low mass stars

In Phan-Bao et al. (2005) we reported the discovery of a new wide binary system of very low mass stars with projected separation of 33 AU. The calibration of the PC3 index to spectral type (Martín et al. 1999b) gives spectral types of M5.5 for component A and M7.5 for component B, with an uncertainty of  $\pm 0.5$  subclass. Figure 1 shows that there is no background star at the position of the system in either the SERC-I image (*I* band) or the DENIS-I image, and therefore demonstrates that the system LP 714-37 is a physical binary. The properties of this system are given in Table 1.

Using the PC3 index to magnitudes relation given in Crifo et al. (2005) and comparison with the DENIS apparent magnitudes give the distance only at 18 pc for our system, this makes LP 714-37AB the closest wide ultracool binary system (separation  $> 30$  AU) in the field up to now. Table 2 lists all known wide ultracool field binary dwarfs.



**Fig. 1.** Archival images of LP 714-37: SERC-I (left, epoch: 1986.940) and DENIS-I (right, epoch: 2000.896). The cross indicates the position of component B at the 2003.915 epoch of the ESO-New Technology Telescope image. Clearly, with the proper motion of the system ( $\mu_\alpha = -117$  mas/yr and  $\mu_\delta = -382$  mas/yr, Phan-Bao et al. 2003), the SERC-I image would easily separate the two objects if they were not physically associated. The size of each image is  $20 \times 20''$ , and North is up and East to the left.

**Table 1.** Summary of the LP 714-37AB binary system

Stars	PC3	SpT	Mass	Sep.	dist.
(1)	(2)	(3)	( $M_\odot$ )	(AU)	(pc)
(1)	(2)	(3)	(4)	(5)	(6)
LP 714-37A	1.32	M5.5	0.11	$33.1 \pm 4.0$	$18.1 \pm 2.2$
LP 714-37B	1.72	M7.5	0.09		

*Column 1:* NLTT name. *Columns 2 & 3:* The PC3 index and spectral types derived from the (PC3, spectral type) relation of Martín et al. (1999b). *Column 4:* Mass determinations for 1–5 Gyr from the models of Baraffe et al. (1998). *Column 5:* Projected separation. *Column 6:* Spectrophotometric distance.

### 3. Discussion

Recent surveys demonstrate that VLM binaries with large separations ( $> 30$  AU) are rare in the field, but can be found in young associations and clusters: 2MASS J1101-7732 (240 AU separation, in Chamaeleon I, Luhman 2004a); 2MASS J1207-3932 (55 AU, in TW Hydrae, Chauvin et al. 2004). Phan-Bao et al. (2005) reported the discovery of a 33 AU VLM binary, Martín et al. (2000) found that CFHT-PI-18 is a 35 AU VLM binary; and very recently, Billères et al. (2005) discovered an M8.5+L0 pair (DENIS-P J055146.0-443412.2) with a physical separation over 200 AU, all of them in the field.

Since ejection models suggest that the binary DB systems that do exist must be close. And therefore the existence of the extremely wide binaries (e.g., 200 AU separation) becomes a very important test of the ejection models (Bate & Bonnell 2005). One should note however that the numerical models to date suffer from small number statistics. One should also note that the relevant quantity is the total mass of the system, and that the apparent binaries could possibly be triple or higher order multiple systems, with a correspondingly higher total mass. This would make them analogs of the GJ1245ABC triple system, which consists of two M5.5 and

**Table 2.** Widely separated binary systems (separations  $> 30$  AU) in the field

Stars	Sep.	SpT <sub>A</sub> /SpT <sub>B</sub>	Dist.	Ref.
(1)	(AU)	(3)	(pc)	(5)
(1)	(2)	(3)	(4)	(5)
LP 714-37	33	M5.5/M7.5	18	1
CFHT-PI-18	35	M8.0/M8.0	105	2,3
DENIS-P J0551-4434	200	M8.5/L0.0	100	4

*Column 1:* Star name. *Columns 2 & 3:* Projected separation and spectral types of components. *Column 4:* Distance estimate. *Column 5:* References: (1) Phan-Bao et al. (2005); (2) Martín et al. (2000); (3) Bouy et al. (2003); (4) Billères et al. (2005).

one  $\sim$ M8 dwarfs with separations of 32 and 5 AU. Triple systems could thus potentially explain the apparent excess of wide VLM binaries, and adaptive optics imaging of LP 714-37, DENIS-P J055146.0-443412.2 would be of obvious interest to clarify its true multiplicity.

*Acknowledgements.* P-B.N. is grateful to the DENIS consortium for access to the DENIS data used by his very low mass stars search. Partial funding was provided by NSF grant AST 02-05862. P-B.N. would like to thank the LOC of the La Palma workshop on Ultra-Low Mass Star Formation for organizing a very enjoyable conference.

### References

- Baraffe, I., Chabrier, G., et al: 1998, A&A 337, 403  
 Bate, M.R., Bonnell, I.A., & Bromm, V.: 2002, MNRAS 332, L65  
 Bate, M.R., & Bonnell, I.A.: 2005, MNRAS 356, 1201  
 Billères, M., Delfosse, X., Beuzit, J.-L., et al.: 2005, A&A 440, L55  
 Bouy, H., Brandner, W., Martín, E.L., et al.: 2003, AJ 126, 1526  
 Burgasser, A.J., Kirkpatrick, J.D., et al.: 2003, ApJ 586, 512  
 Chauvin, G., Lagrange, A.-M., et al.: 2004, A&A 425, L29  
 Close, L.M., Siegler, N., et al.: 2002, ApJ 567, L53  
 Close, L.M., Siegler, N., et al.: 2003, ApJ 587, 407  
 Crifo, F., Phan-Bao, N., Delfosse, X., et al.: 2005, accepted by A&A  
 Forveille, T., Beuzit, J.-L., Delorme, P., et al. 2005, A&A, L435  
 Kroupa, P., Bouvier, J., et al.: 2003, MNRAS 346, 354  
 Luhman, K.L.: 2004a, ApJ 614, 398  
 Luhman, K.L.: 2004b, ApJ 617, 1216  
 Martín, E. L., Brandner, W., & Basri, G.: 1999a, Sci 283, 1718  
 Martín, E. L., Delfosse, X., et al.: 1999b, AJ 118, 2466  
 Martín, E. L., Brandner, W., Bouvier, J., et al.: 2000, ApJ 543, 299  
 Martín, E. L., Barrado y Navascués, D., et al.: 2003, ApJ 594, 525  
 Phan-Bao, N., Crifo, F., Delfosse, X., et al.: 2003, A&A 401, 959  
 Phan-Bao, N., Martín, E., Reylé, C., et al.: 2005, A&A 439, L19  
 Reipurth, B., & Clarke, C.: 2001, AJ 122, 432